Universal Patient Care

Scene Safe

YES

NO

Call for help / additional resources
Stage until scene safe

Bring all necessary equipment to patient
Demonstrate professionalism and courtesy
Mass assembly consider WMD

Utilize appropriate PPE

Consider Airborne, Contact, or Droplet Isolation ***if indicated***

Initial assessment
BLS maneuvers
Initiate oxygen ***if indicated***

Adult Assessment Procedure

Pediatric Assessment Procedure
Use Broselow-Luten tape

Required VS:
Blood pressure
Palpated pulse rate
Respiratory rate
Pulse ox if available

If Indicated:
Glucose
12 Lead ECG
Temperature
Pain scale
CO Monitoring
EtCO2 Monitoring

Medical or Trauma

Trauma Patient

Medical Patient

Evaluate MOI

Mental Status Exam

Significant MOI

No Significant MOI

Unresponsive

Responsive

Primary and Secondary trauma assessment

Primary and Secondary trauma assessment

Focused assessment on specific injury

Primary and secondary assessment

Obtain history of present illness from available sources / scene survey

Obtain Chief Complaint

Primary and Secondary assessment
Focused assessment on specific complaint

Spinal Motion Restriction Procedure / Protocol TB 8 ***as indicated***

Obtain VS

Obtain SAMPLE

Repeat assessment while preparing for transport

Exit to
Age Appropriate Protocol(s)
as indicated

Continue on-going assessment
Repeat initial VS
Evaluate interventions / procedures

Transfer
Patient hand-off includes patient information, personal property and summary of care and response to care

Patient does not fit specific protocol

Notify Destination or Contact Medical Control

Universal Protocol Section

Universal Patient Care

Universal Protocol Section

- **Pearls**
- **Recommended Exam: Minimal exam if not noted on the specific protocol is vital signs, mental status with GCS, and location of injury or complaint.**
- **Any patient contact which does not result in an EMS transport must have a completed disposition form.**
- **Vital signs should be obtained before, 10 minutes after, and at patient hand off with all pain medications.**
- **2 complete vital sign acquisitions should occur at a minimum with a patient encounter.**
- **Patient Refusal**

  Patient refusal is a high risk situation. Encourage patient to accept transport to medical facility.
  Encourage patient to allow an assessment, including vital signs. Documentation of the event is very important including a mental status assessment describing the patient's capacity to refuse care.
  Guide to Assessing capacity:
  C:***Patient should be able to communicate a clear choice:*** This should remain stable over time. Inability to communicate a choice or an inability to express the choice consistently demonstrates incapacity.
  R: ***Relevant information is understood:*** Patient should be able to display a factual understanding of the illness, the options and risks and benefits.
  A: ***Appreciation of the situation:*** Ability to communicate an understanding of the facts of the situation. They should be able to recognize the significance of the outcome potentially from their decision.
  M: ***Manipulation of information in a rational manner:*** Demonstrate a rational process to come to a decision. Should be able to describe the logic they are using to come to the decision, though you may not agree with decision.
- **Pediatric Patient General Considerations:**

  A pediatric patient is defined by fitting a Length-based Resuscitation Tape, Age ≤ 15, weight ≤ 49 kg.
  Patients off the Broselow-Luten tape should have weight based medications until age ≥ 16 or weight ≥ 50 kg.
  Special needs children may require continued use of Pediatric based protocols regardless of age and weight.
  Initial assessment should utilize the Pediatric Assessment Triangle which encompasses Appearance, Work of Breathing and Circulation to skin.
  The order of assessment may require alteration dependent on the developmental state of the pediatric patient. Generally the child or infant should not be separated from the caregiver unless absolutely necessary during assessment and treatment.
- Timing of transport should be based on patient's clinical condition and the transport policy.
- Never hesitate to contact medical control for patient who refuses transport.
- Blood Pressure is defined as a Systolic / Diastolic reading. A palpated Systolic reading may be necessary at times.
- SAMPLE: Signs / Symptoms; Allergies; Medications; PMH; Last oral intake; Events leading to illness / injury

Triage



Universal Protocol Section

Triage

Universal Protocol Section

Pearls

- .**When approaching a multiple casualty incident where resources are limited:**
  Triage decisions must be made rapidly with less time to gather information
  Emphasis shifts from ensuring the best possible outcome for an individual patient to ensuring the best possible outcome for the greatest number of patients.
- **Scene Size Up:**
  1. **Conduct a scene size up. Assure well being of responders. Determine or ensure scene safety before entering. If there are several patients with the same complaints consider HazMat, WMC or CO poisoning.**
  2. **Take Triage system kit.**
  3. **Determine number of patients. Communicate the number of patients and nature of the incident, establish command and establish a medical officer and triage officer if personnel available**
- **Triage is a continual process and should recur in each section as resources allow.**
- **Step 1: Global sorting:**
  Call out to those involved in the incident to walk to a designated area and assess third.
  For those who cannot walk, have them wave / indicate a purposeful movement and assess them second.
  Those involved who are not moving or have an obvious life threat, assess first.
- **Step 2: Individual assessments:**
  Control major hemorrhage
  Open airway and if child, give 2 rescue breaths
  Perform Needle Chest Decompression Procedure if indicated.
  Administer injector antidotes if indicated
- **Assess the first patient you encounter using the three objective criteria which can be remembered by RPM.**
  R: Respiratory
  P: Perfusion
  M: Mental Status
- If your patient falls into the RED TAG category, stop, place RED TAG and move on to next patient. Attempt only to correct airway problems, treat uncontrolled bleeding, or administer an antidote before moving to next patient.
- **Treatment:**
  Once casualties are triaged focus on treatment can begin. You may need to move patients to treatment areas.
  RED TAGs are moved / treated first followed by YELLOW TAGs. BLACK TAGs should remain in place.
  You may also indicate deceased patients by pulling their shirt / clothing over their head.
  As more help arrives then the triage / treatment process may proceed simultaneously.
- Capillary refill can be altered by many factors including skin temperature. Age-appropriate heart rate may also be used in triage decisions.
- SMART triage tag system is utilized in NC.

Abdominal Pain
Vomiting and Diarrhea

History
- Age
- Time of last meal
- Last bowel movement/emesis
- Improvement or worsening with food or activity
- Duration of problem
- Other sick contacts
- Past medical history
- Past surgical history
- Medications
- Menstrual history (pregnancy)
- Travel history
- Bloody emesis / diarrhea

Signs and Symptoms
- Pain
- Character of pain (constant, intermittent, sharp, dull, etc.)
- Distention
- Constipation
- Diarrhea
- Anorexia
- Radiation

Associated symptoms:
Fever, headache, blurred vision, weakness, malaise, myalgias, cough, headache, dysuria, mental status changes, rash

Differential
- CNS (increased pressure, headache, stroke, CNS lesions, trauma or hemorrhage, vestibular)
- Myocardial infarction
- Drugs (NSAID's, antibiotics, narcotics, chemotherapy)
- GI or Renal disorders
- Diabetic ketoacidosis
- OB-Gyn disease (ovarian cyst, PID, Pregnancy)
- Infections (pneumonia, influenza)
- Electrolyte abnormalities
- Food or toxin induced
- Medication or Substance abuse
- Psychological

	Consider Blood Glucose Analysis Procedure
B	12 Lead ECG Procedure
A	IV Procedure
P	Cardiac Montior
	Age Appropriate Diabetic Protocol AM 2 / PM 2 ***if indicated***
	Pain Control Protocol UP 11 ***if indicated***
	Age Appropriate Cardiac Protocol(s) ***if indicated***

Serious Signs / Symptoms
Hypotension, poor perfusion, shock

YES

A	IV Procedure **P** IO Procedure
A	Consider 2 Large Bore Sites
A	**Normal Saline 500 mL Bolus** Repeat as needed Titrate SPB ≥ 90 mmHg **Maximum 2 L** **Peds: 20 mL/kg IV / IO** Repeat as needed Titrate to Age Appropriate SBP ≥ 70 + 2 x Age **Maximum 60 mL/kg**
P	**Ondansetron 4 mg IV / IO / IM** **Peds: 0.15 mg/kg IV / IO Max of 4 mg**
	Age Appropriate Hypotension / Shock Protocol AM 5 / PM 3 ***if indicated***

NO

A	**Normal Saline IV TKO Or Saline Lock**
P	**Ondansetron 4 mg IV / IO / IM** **Peds: 0.15 mg/kg IV / IO Max of 4mg** **Promethazine 12.5 mg IV/ IO / IM** ½ Dose if > 60 y/o May repeat x 1 as needed

Monitor and Reassess

Notify Destination or Contact Medical Control

Universal Protocol Section

Abdominal Pain
Vomiting and Diarrhea

Universal Protocol Section

Pearls

- **Recommended Exam: Mental Status, Skin, HEENT, Neck, Heart, Lungs, Abdomen, Back, Extremities, Neuro**
- **Age specific blood pressure 0 – 28 days > 60 mmHg, 1 month - 1 year > 70 mmHg, 1 - 10 years > 70 + (2 x age) mmHg and 11 years and older > 90 mmHg.**
- **Abdominal / back pain in women of childbearing age should be treated as pregnancy related until proven otherwise.**
- **The diagnosis of abdominal aneurysm should be considered with abdominal pain, with or without back and / or lower extremity pain or diminished pulses, especially in patients over 50 and / or patients with shock/ poor perfusion. Notify receiving facility early with suspected abdominal aneurysm.**
- **Consider cardiac etiology in patients > 50, diabetics and / or women especially with upper abdominal complaints.**
- **Repeat vital signs after each fluid bolus.**
- **Heart Rate: One of the first clinical signs of dehydration, almost always increased heart rate, tachycardia increases as dehydration becomes more severe, very unlikely to be significantly dehydrated if heart rate is close to normal.**
- **Promethazine (Phenergan) may cause sedative effects in pediatric patients and ages ≥ 60 and the debilitated, etc.) When giving promethazine IV dilute with 10 mL of normal saline and administer slowly as it can also harm the veins.**
- Beware of vomiting only in children. Pyloric stenosis, bowel obstruction, and CNS processes (bleeding, tumors, or increased CSF pressures) all often present with vomiting.
- Document the mental status and vital signs prior to administration of Promethazine (Phenergan).
- Isolated vomiting may be caused by pyloric stenosis, bowel obstruction, and CNS processes (bleeding, tumors, or increased CSF pressures).
- Vomiting and diarrhea are common symptoms, but can be the symptoms of uncommon and serious pathology such as stroke, carbon monoxide poisoning, acute MI, new onset diabetes, diabetic ketoacidosis (DKA), and organophosphate poisoning. Maintain a high index of suspicion.

Altered Mental Status

History
- Known diabetic, medic alert tag
- Drugs, drug paraphernalia
- Report of illicit drug use or toxic ingestion
- Past medical history
- Medications
- History of trauma
- Change in condition
- Changes in feeding or sleep habits

Signs and Symptoms
- Decreased mental status or lethargy
- Change in baseline mental status
- Bizarre behavior
- Hypoglycemia (cool, diaphoretic skin)
- Hyperglycemia (warm, dry skin; fruity breath; Kussmaul respirations; signs of dehydration)
- Irritability

Differential
- Head trauma
- CNS (stroke, tumor, seizure, infection)
- Cardiac (MI, CHF)
- Hypothermia
- Infection (CNS and other)
- Thyroid (hyper / hypo)
- Shock (septic, metabolic, traumatic)
- Diabetes (hyper / hypoglycemia)
- Toxicological or Ingestion
- Acidosis / Alkalosis
- Environmental exposure
- Pulmonary (Hypoxia)
- Electrolyte abnormality
- Psychiatric disorder

	Age Appropriate Airway Protocol(s) AR 1, 2, 3, 5, 6 ***if indicated***
	Blood Glucose Analysis Procedure
B	12 Lead ECG Procedure
A	IV Procedure / P IO Procedure
	Age Appropriate Diabetic Protocol(s) AM 2 / PM 2 ***if indicated***

Signs of shock / Poor perfusion Traumatic Injury — YES → Exit to Age Appropriate Hypotension / Shock Protocol AM 5 / PM 3, Multiple Trauma Protocol TB 6, Head Injury Protocol TB 5

NO

Signs of OD / Toxicology — YES → Exit to Overdose / Toxic Exposure Protocol TE 7

NO

Signs of CVA Or Seizure — YES → Exit to Suspected Stroke Protocol UP 14, Seizure Protocol UP 13

NO

Signs of Hypo / Hyperthermia — YES → Exit to Hypothermia Protocol TE 5, Hyperthermia Protocol TE 4

NO

Arrhythmia / STEMI / CP — YES → Exit to Age Appopriate Appropriate Cardiac Protocol(s)

NO

Fever / Sepsis — YES → Exit to Fever Protocol UP 10, Suspected Sepsis Protocol UP 15

NO → **Notify Destination or Contact Medical Control**

Universal Protocol Section

Altered Mental Status

Universal Protocol Section

Pearls

- **Recommended Exam: Mental Status, HEENT, Skin, Heart, Lungs, Abdomen, Back, Extremities, Neuro.**
- **AMS may present as a sign of an environmental toxin or Haz-Mat exposure - protect personal safety.**
- **General:**
  - **The patient with AMS poses one of the most significant challenges.**
  - **A careful assessment of the patient, the scene and the circumstances should be undertaken.**
  - **Assume the patient has a life threatening cause of their AMS until proven otherwise.**
  - **Pay careful attention to the head exam for signs of bruising or other injury.**
  - **Information found at the scene must be communicated to the receiving facility.**
- **Substance misuse:**
  - Patients ingesting substances can pose a great challenge.
  - DO NOT assume recreational drug use and / or alcohol are the sole reasons for AMS.
  - Misuse of alcohol may lead to hypoglycemia.
  - More serious underlying medical and trauma conditions may be the cause.
- **Behavioral health:**
  - The behavioral health patient may present a great challenge in forming a differential.
  - DO NOT assume AMS is the result solely of an underlying psychiatric etiology.
  - Often an underlying medial or trauma condition precipitates a deterioration of a patients underlying disease.
- **Spinal Motion Restriction / Trauma:**
  - Only utilize spinal immobilization if the situation warrants.
  - The patient with AMS may worsen with increased agitation when immobilized.
- **It is safer to assume hypoglycemia than hyperglycemia if doubt exists. Recheck blood glucose after Dextrose or Glucagon**
- Consider Restraints if necessary for patient's and/or personnel's protection per the restraint procedure.

Back Pain

History
- Age
- Past medical history
- Past surgical history
- Medications
- Onset of pain / injury
- Previous back injury
- Traumatic mechanism
- Location of pain
- Fever
- Improvement or worsening with activity

Signs and Symptoms
- Pain (paraspinous, spinous process)
- Swelling
- Pain with range of motion
- Extremity weakness
- Extremity numbness
- Shooting pain into an extremity
- Bowel / bladder dysfunction

Differential
- Muscle spasm / strain
- Herniated disc with nerve compression
- Sciatica
- Spine fracture
- Kidney stone
- Pyelonephritis
- Aneurysm
- Pneumonia
- Spinal Epidural Abscess
- Metastatic Cancer
- AAA



Universal Protocol Section

Back Pain

Universal Protocol Section

Pearls

- **Recommended Exam: Mental Status, Heart, Lungs, Abdomen, Neuro, Lower extremity perfusion**
- **Back pain is one of the most common complaints in medicine and effects more than 90 % of adults at some point in their life. Back pain is also common in the pediatric population. Most often it is a benign process but in some circumstances can be life or limb threatening.**
- **Consider pregnancy or ectopic pregnancy with abdominal or back pain in women of childbearing age.**
- **Consider abdominal aortic aneurysm with abdominal pain especially in patients over 50 and/or patients with shock/ poor perfusion. Patients may have abdominal pain and / or lower extremity pain with diminished pulses, . Notify receiving facility early with suspected abdominal aneurysm.**
- **Consider cardiac etiology in patients > 50, diabetics and / or women especially with upper abdominal complaints.**
- **Red Flags which may signal more serious process associated with back pain:**
  - Age > 50 or < 18
  - Neurological deficit (leg weakness, urinary retention, or bowel incontinence)
  - IV Drug use
  - Fever
  - History of cancer, either current or remote
  - Night time pain in pediatric patients
- **Cauda equina syndrome is where the terminal nerves of spinal cord are being compressed (Symptoms include):.**
  - Saddle anesthesia
  - Recent onset of bladder and bowel dysfunction. (Urine retention and bowel incontinence)
  - Severe or progressive neurological deficit in the lower extremity.
  - Motor weakness of thigh muscles or foot drop
- **Back pain associated with infection:**
  - Fever / chills.
  - IV Drug user (consider spinal epidural abscess)
  - Recent bacterial infection like pneumonia.
  - Immune suppression such as HIV or patients on chronic steroids like prednisone.
  - Meningitis.
- **Spinal motion restriction in patients with underlying spinal deformity should be maintained in their functional position.**
- Kidney stones typically present with an acute onset of flank pain which radiates around to the groin area.

Behavioral

History
- Situational crisis
- Psychiatric illness/medications
- Injury to self or threats to others
- Medic alert tag
- Substance abuse / overdose
- Diabetes

Signs and Symptoms
- Anxiety, agitation, confusion
- Affect change, hallucinations
- Delusional thoughts, bizarre behavior
- Combative violent
- Expression of suicidal / homicidal thoughts

Differential
- Altered Mental Status differential
- Alcohol Intoxication
- Toxin / Substance abuse
- Medication effect / overdose
- Withdrawal syndromes
- Depression
- Bipolar (manic-depressive)
- Schizophrenia
- Anxiety disorders

Call for help / additional resources
Stage until scene safe

Blood Glucose Analysis Procedure ***if indicated***

Age Appropriate Diabetic Protocol AM 3 / PM 2 if indicated

Altered Mental Status Protocol UP 4
Overdose / Toxic Ingestion Protocol TE 7 if indicated

Head Trauma Protocol TB 5
Multiple Trauma Protocol TB 6 if indicated

Excited Delirium Syndrome
Paranoia, disorientation, hyper-aggression, hallucination, tachycardia, increased strength, hyperthermia

YES

P
Midazolam 2.5 mg IV / IO
5 mg IM
Age ≥ 65
1 – 2.5 mg IV / IO
2.5 mg IM
Peds: 0.1 – 0.2 mg/kg
IV / IO / IM
Repeat every 2-3 minutes as needed

NO

Aggressive, Violent, Agitation

Threat to Self / others
Setting of Psychosis

YES

Consider Restraint Physical Procedure

Monitor per restraint procedure ***if indicated***

A IV Procedure P IO Procedure

P
Age ≥ 12
Haloperidol 2.5 - 5 mg IM / IV
Age ≥ 65
2.5 mg IM / IV
May repeat every 5 as needed
Maximum 10 mg

P
Midazolam 2.5 mg IV / IO
5 mg IM
Age ≥ 65
1 – 2.5 mg IV / IO
2.5 mg IM
Peds: 0.1 – 0.2 mg/kg
IV / IO / IM
Repeat every 2-3 minutes as needed

A IV Procedure P IO Procedure

A Consider 2 Large Bore Sites

A
Normal Saline 1 L Bolus
Then **150 – 200 mL / hr**
May repeat **500 mL** Bolus as needed
Maximum 2 L
Peds: 20 – 60 mL/kg IV / IO
Maximum 60 mL/kg

External Cooling Measures

Consider Restraint Physical Procedure

Monitor per restraint procedure ***if indicated***

P Cardiac Monitor

NO

Monitor and Reassess

Notify Destination or Contact Medical Control

Universal Protocol Section

UP 6 (ROBESON)

This protocol has been altered from the original NCCEP Protocol by the local EMS Medical Director

Behavioral

Universal Protocol Section

Pearls

- **Recommended Exam: Mental Status, Skin, Heart, Lungs, Neuro**
- **Crew / responders safety is the main priority.**
- **Any patient who is handcuffed or restrained by Law Enforcement and transported by EMS must be accompanied by law enforcement in the ambulance.**
- **Consider Haldol for patients with history of psychosis or a benzodiazepine for patients with presumed substance abuse.**
- **Haldol is acceptable treatment in pediatric patients ≥ 12 years old. Safety and efficacy is not established in younger ages.**
- **All patients who receive either physical or chemical restraint must be continuously observed by ALS personnel on scene or immediately upon their arrival.**
- Be sure to consider all possible medical/trauma causes for behavior (hypoglycemia, overdose, substance abuse, hypoxia, head injury, etc.)
- Do not irritate the patient with a prolonged exam.
- Do not overlook the possibility of associated domestic violence, child, or geriatric abuse.
- Do not position or transport any restrained patient is such a way that could impact the patients respiratory or circulatory status.
- **Excited Delirium Syndrome:**

  Medical emergency: Combination of delirium, psychomotor agitation, anxiety, hallucinations, speech disturbances, disorientation, violent / bizarre behavior, insensitivity to pain, hyperthermia and increased strength. Potentially life-threatening and associated with use of physical control measures, including physical restraints and Tasers. Most commonly seen in male subjects with a history of serious mental illness and/or acute or chronic drug abuse, particularly stimulant drugs such as cocaine, crack cocaine, methamphetamine, amphetamines or similar agents. Alcohol withdrawal or head trauma may also contribute to the condition.
- **If patient is suspected of EDS suffers cardiac arrest, consider a fluid bolus and sodium bicarbonate early**
- **Extrapyramidal reactions:**

  Condition causing involuntary muscle movements or spasms typically of the face, neck and upper extremities. May present with contorted neck and trunk with difficult motor movements. Typically an adverse reaction to antipsychotic drugs like Haloperidol and may occur with your administration. When recognized give **Diphenhydramine 50 mg IV / IO / IM /** in adults or **1 mg/kg IV / IO / IM** in pediatrics.

Dental Problems

History
- Age
- Past medical history
- Medications
- Onset of pain / injury
- Trauma with "knocked out" tooth
- Location of tooth
- Whole vs. partial tooth injury

Signs and Symptoms
- Bleeding
- Pain
- Fever
- Swelling
- Tooth missing or fractured

Differential
- Decay
- Infection
- Fracture
- Avulsion
- Abscess
- Facial cellulitis
- Impacted tooth (wisdom)
- TMJ syndrome
- Myocardial infarction



Universal Protocol Section

Pearls
- **Recommended Exam: Mental Status, HEENT, Neck, Chest, Lungs, Neuro**
- Significant soft tissue swelling to the face or oral cavity can represent a cellulitis or abscess.
- Scene and transport times should be minimized in complete tooth avulsions. Reimplantation is possible within 4 hours if the tooth is properly cared for.
- Occasionally cardiac chest pain can radiate to the jaw.
- All pain associated with teeth should be associated with a tooth which is tender to tapping or touch (or sensitivity to cold or hot).

Emergencies Involving Indwelling Central Lines

History

- Central Venous Catheter Type
  - Tunneled Catheter (Broviac / Hickman)
  - PICC (peripherally inserted central catheter
  - Implanted catheter (Mediport / Hickman)
- Occlusion of line
- Complete or partial dislodge
- Complete or partial disruption

Signs and Symptoms

- External catheter dislodgement
- Complete catheter dislodgement
- Damaged catheter
- Bleeding at catheter site
- Internal bleeding
- Blood clot
- Air embolus
- Erythema, warmth or drainage about catheter site indicating infection

Differential

- Fever
- Hemorrhage
- Reactions from home nutrient or medication
- Respiratory distress
- Shock



Universal Protocol Section

Pearls

- **Always talk to family / caregivers as they have specific knowledge and skills.**
- **Use strict sterile technique when accessing / manipulating an indwelling catheter.**
- **Cardiac arrest: May access central catheter and utilize if functioning properly.**
- Do not attempt to force catheter open if occlusion evident.
- Some infusions may be detrimental to stop. Ask family or caregiver if it is appropriate to stop or change infusion.
- Hyperalimentation infusions (IV nutrition): If stopped for any reason monitor for hypoglycemia.

Epistaxis

History
- Age
- Past medical history
- Medications (HTN, anticoagulants, aspirin, NSAIDs)
- Previous episodes of epistaxis
- Trauma
- Duration of bleeding
- Quantity of bleeding

Signs and Symptoms
- Bleeding from nasal passage
- Pain
- Nausea
- Vomiting

Differential
- Trauma
- Infection (viral URI or Sinusitis)
- Allergic rhinitis
- Lesions (polyps, ulcers)
- Hypertension



Universal Protocol Section

Pearls
- **Recommended Exam: Mental Status, HEENT, Heart, Lungs, Neuro**
- **Age specific hypotension: 0 – 28 days < 60 mmHg, 1 month – 1 year < 70 mmHg, 1 year – 10 years < 70 + (2 x age) mmHg, 11 years and greater < 90 mmHg.**
- It is very difficult to quantify the amount of blood loss with epistaxis.
- Bleeding may also be occurring posteriorly. Evaluate for posterior blood loss by examining the posterior pharnyx.
- Anticoagulants include warfarin (Coumadin), Apixaban (Elequis), heparin, enoxaparin (Lovenox), dabigatran (Pradaxa), rivaroxaban (Xarelto), and many over the counter headache relief powders.
- Anti-platelet agents like aspirin, clopidogrel (Plavix), aspirin/dipyridamole (Aggrenox), and ticlopidine (Ticlid) can contribute to bleeding.

Fever / Infection Control

History
- Age
- Duration of fever
- Severity of fever
- Past medical history
- Medications
- Immunocompromised (transplant, HIV, diabetes, cancer)
- Environmental exposure
- Last acetaminophen or ibuprofen

Signs and Symptoms
- Warm
- Flushed
- Sweaty
- Chills/Rigors

Associated Symptoms (Helpful to localize source)
- myalgias, cough, chest pain, headache, dysuria, abdominal pain, mental status changes, rash

Differential
- Infections / Sepsis
- Cancer / Tumors / Lymphomas
- Medication or drug reaction
- Connective tissue disease
  - Arthritis
  - Vasculitis
- Hyperthyroidism
- Heat Stroke
- Meningitis



Universal Protocol Section

Pearls
- **Recommended Exam: Mental Status, Skin, HEENT, Neck, Heart, Lungs, Abdomen, Back, Extremities, Neuro**
- Febrile seizures are more likely in children with a history of febrile seizures and with a rapid elevation in temperature.
- Patients with a history of liver failure should not receive acetaminophen.
- **Droplet precautions** include standard PPE plus a standard surgical mask for providers who accompany patients in the back of the ambulance and a surgical mask or NRB O2 mask for the patient. This level of precaution should be utilized when influenza, meningitis, mumps, streptococcal pharyngitis, and other illnesses spread via large particle droplets are suspected. A patient with a potentially infectious rash should be treated with droplet precautions.
- **Airborne precautions** include standard PPE plus utilization of a gown, change of gloves after every patient contact, and strict hand washing precautions. This level of precaution is utilized when multi-drug resistant organisms (e.g. MRSA), scabies, or zoster (shingles), or other illnesses spread by contact are suspected.
- **All-hazards precautions** include standard PPE plus airborne precautions plus contact precautions. This level of precaution is utilized during the initial phases of an outbreak when the etiology of the infection is unknown or when the causative agent is found to be highly contagious (e.g. SARS).
- Rehydration with fluids increases the patient's ability to sweat and improves heat loss.
- All patients should have drug allergies documented prior to administering pain medications.
- Allergies to NSAIDs (non-steroidal anti-inflammatory medications) are a contraindication to Ibuprofen. Do not give to patients who have renal disease or renal transplant.
- NSAIDs should not be used in the setting of environmental heat emergencies.
- **Do not** give aspirin to a child, age ≤ 15 years.
- Agency Medical Director may require contact of medical control prior to EMT / EMR administering any medication.

Pain Control

History
- Age
- Location
- Duration
- Severity (1 - 10)
- If child use Wong-Baker faces scale
- Past medical history
- Medications
- Drug allergies

Signs and Symptoms
- Severity (pain scale)
- Quality (sharp, dull, etc.)
- Radiation
- Relation to movement, respiration
- Increased with palpation of area

Differential
- Per the specific protocol
- Musculoskeletal
- Visceral (abdominal)
- Cardiac
- Pleural / Respiratory
- Neurogenic
- Renal (colic)

Enter from
Protocol based on **Specific Complaint**

Assess Pain Severity

Use combination of Pain Scale, Circumstances, MOI, Injury or Illness severity

Mild

B

Ibuprofen 10 mg/kg PO (Ped)
400 - 800 mg (Adult)
Or
Acetaminophen 15 mg/kg (Ped)
500 - 1000 mg (Adult)
Or
Aspirin 324 to 650 mg PO
(≥ 16 only)

A

Consider IV Procedure
if indicated

Moderate to Severe

A

IV Procedure | **P** | IO Procedure

Ketorolac 30 mg IV / IO
60 mg IM
Maximum 60 mg
Peds: 0.5 mg/kg IV / IO / IM
Maximum 30 mg

P

Cardiac Monitor

Fentanyl 50 – 75 mcg IV / IO
Repeat **25 mcg** every 20 minutes
Maximum 150 mcg
Peds: 1 mcg/kg IV / IO / IM
May repeat 0.5 mg/kg every 5 minutes
Maximum 2 mcg/kg
Or
Morphine 4 mg IV / IO / IM
Repeat **2 mg** every 5 minutes as needed
Peds: 0.1 mg/kg IV / IO / IM
May repeat every 5 minutes
Maximum 10 mg

Monitor and Reassess
Every 10 minutes following sedative

Monitor and Reassess

Notify Destination or
Contact Medical Control

Universal Protocol Section

Pain Control

Universal Protocol Section

Pearls

- **Recommended Exam: Mental Status, Area of Pain, Neuro**
- **Pain severity (0-10) is a vital sign to be recorded before and after PO, IV, IO or IM medication delivery and at patient hand off. Monitor BP closely as sedative and pain control agents may cause hypotension.**
- **Both arms of the treatment may be used in concert. For patients in Moderate pain for instance, you may use the combination of an oral medication and parenteral if no contraindications are present.**
- **Pediatrics:**
  For children use Wong-Baker faces scale or the FLACC score (see Assessment Pain Procedure)
  Use Numeric (> 9 yrs), Wong-Baker faces (4-16yrs) or FLACC scale (0-7 yrs) as needed to assess pain
- **Vital signs should be obtained before, 10 minutes after, and at patient hand off with all pain medications.**
- All patients who receive IM or IV medications must be observed 15 minutes for drug reaction in the event no transport occurs.
- Do not administer any PO medications for patients who may need surgical intervention such as open fractures or fracture deformities, headaches, or abdominal pain.
- **Ketorolac (Toradol) and Ibuprofen should not be used in patients with known renal disease or renal transplant, in patients who have known drug allergies to NSAID's (non-steroidal anti-inflammatory medications), with active bleeding, headaches, abdominal pain, stomach ulcers or in patients who may need surgical intervention such as open fractures or fracture deformities.**
- Do not administer **Acetaminophen** to patients with a history of liver disease.
- Burn patients may required higher than usual opioid doses to titrate adequate pain control.
- Consider agency-specific anti-emetic(s) for nausea and/or vomiting.

Police Custody

History
- Traumatic Injury
- Drug Abuse
- Cardiac History
- History of Asthma
- Psychiatric History

Signs and Symptoms
- External signs of trauma
- Palpitations
- Shortness of breath
- Wheezing
- Altered Mental Status
- Intoxication/Substance Abuse

Differential
- Agitated Delirium Secondary to Psychiatric Illness
- Agitated Delirium Secondary to Substance Abuse
- Traumatic Injury
- Closed Head Injury
- Asthma Exacerbation
- Cardiac Dysrhythmia



Universal Protocol Section

Police Custody

Universal Protocol Section

Pearls

- **Patient does not have to be in police custody or under arrest to utilize this protocol.**
- **Local EMS agencies should formulate a policy with local law enforcement agencies concerning patients requiring EMS and Law Enforcement simultaneously. Agencies should work together to formulate a disposition in the best interest of the patient.**
- **Patients restrained by law enforcement devices must be transported accompanied by a law enforcement officer in the patient compartment who is capable of removing the devices. However when rescuers have utilized restraints in accordance with Restraint Procedure, the law enforcement agent may follow behind the ambulance during transport.**
- **All patients who receive either physical or chemical restraint must be continuously observed by ALS personnel on scene or immediately upon their arrival.**
- The responsibility for patient care rests with the highest authorized medical provider on scene per North Carolina law.
- If an asthmatic patient is exposed to pepper spray and released to law enforcement, all parties should be advised to immediately contact EMS if wheezing/difficulty breathing occurs.
- All patients with decision-making capacity in police custody retain the right to participate in decision making regarding their care and may request care or refuse care of EMS.
- If extremity / chemical / law enforcement restraints are applied, follow Restraint Procedure.
- **Consider Haldol for patients with history of psychosis or a benzodiazepine for patients with presumed substance abuse.**
- **Haldol is acceptable treatment in pediatric patients ≥ 12 years old. Safety and efficacy is not established in younger ages.**
- **<u>Excited Delirium Syndrome:</u>**

  Medical emergency: Combination of delirium, psychomotor agitation, anxiety, hallucinations, speech disturbances, disorientation, violent / bizarre behavior, insensitivity to pain, hyperthermia and increased strength. Potentially life-threatening and associated with use of physical control measures, including physical restraints and Tasers.
  Most commonly seen in male subjects with a history of serious mental illness and/or acute or chronic drug abuse, particularly stimulant drugs such as cocaine, crack cocaine, methamphetamine, amphetamines or similar agents. Alcohol withdrawal or head trauma may also contribute to the condition.
  If patient suspected of EDS suffers cardiac arrest, consider a fluid bolus and sodium bicarbonate early.
- Do not position or transport any restrained patient is such a way that could impact the patients respiratory or circulatory status.
- Patients exposed to chemical spray, with or without history of respiratory disease, may develop respiratory complaints up to 20 minutes post exposure.

Seizure

History
- Reported / witnessed seizure activity
- Previous seizure history
- Medical alert tag information
- Seizure medications
- History of trauma
- History of diabetes
- History of pregnancy
- Time of seizure onset
- Document number of seizures
- Alcohol use, abuse or abrupt cessation
- Fever

Signs and Symptoms
- Decreased mental status
- Sleepiness
- Incontinence
- Observed seizure activity
- Evidence of trauma
- Unconscious

Differential
- CNS (Head) trauma
- Tumor
- Metabolic, Hepatic, or Renal failure
- Hypoxia
- Electrolyte abnormality (Na, Ca, Mg)
- Drugs, Medications, Non-compliance
- Infection / Fever
- Alcohol withdrawal
- Eclampsia
- Stroke
- Hyperthermia
- Hypoglycemia

Loosen any constrictive clothing
Protect patient

Blood Glucose Analysis Procedure

Altered Mental Status
Protocol UP 4
if indicated

Behavioral Protocol UP 6
if indicated

A IV Procedure **If indicated** | **P** IO Procedure **If indicated**

P Cardiac Monitor ***if indicated***

Active Seizure Activity

YES →

Age Appropriate
Airway Protocol(s) AR 1, 2, 3, 5, 6
as indicated

Obstetrical Emergency Protocol AO 1
Childbirth Protocol AO 3
if indicated

P

Midazolam 2 – 2.5 mg IV / IO
5 mg IM
Peds: 0.1 – 0.2 mg/kg
IV / IO / IM
May repeat every 3 to 5 minutes as needed
Maxiumum 10 mg
Or
Diazepam 5 mg IV / IO
10 mg Per Rectum
Peds: 0.1 – 0.3 mg/kg IV / IO / IM
Peds rectal: 0.5 mg/kg
Repeat **2 mg** every 3 to 5 minutes as needed
Maximum 10 mg

NO →

Monitor and Reassess

Notify Destination or Contact Medical Control

Universal Protocol Section

Seizure

Universal Protocol Section

Pearls

- **Recommended Exam: Mental Status, HEENT, Heart, Lungs, Extremities, Neuro**
- **Items in Red Text are key performance measures used to evaluate protocol compliance and care**
- **Adult:**
  Midazolam 5 – 10 mg IM is effective in termination of seizures. Do not delay IM administration with difficult IV or IO access. IM Preferred over IO.
- **Pediatrics:**
  Midazolam 0.2 mg/kg (Maximum 10 mg) IM is effective in termination of seizures. Do not delay IM administration with difficult IV or IO access. IM Preferred over IO.
- **Status epilepticus** is defined as two or more successive seizures without a period of consciousness or recovery. This is a true emergency requiring rapid airway control, treatment, and transport.
- **Grand mal seizures (generalized)** are associated with loss of consciousness, incontinence, and tongue trauma.
- **Focal seizures** affect only a part of the body and are not usually associated with a loss of consciousness, but can propagate to generalized seizures with loss of consciousness.
- Be prepared for airway problems and continued seizures.
- Assess possibility of occult trauma and substance abuse.
- In an infant, a seizure may be the only evidence of a closed head injury.
- Be prepared to assist ventilations especially if diazepam or midazolam is used.
- For any seizure in a pregnant patient, follow the OB Emergencies Protocol.
- Diazepam (Valium) is not effective when administered IM. Give IV or Rectally.
- Midazolam is well absorbed when administered IM.

This protocol has been altered from the original NCCEP Protocol by the local EMS Medical Director

Suspected Stroke

History
- Previous CVA, TIA's
- Previous cardiac / vascular surgery
- Associated diseases: diabetes, hypertension, CAD
- Atrial fibrillation
- Medications (blood thinners)
- History of trauma
- Sickle Cell Disease
- Immune disorders
- Congenital heart defects
- Maternal infection / hypertension

Signs and Symptoms
- Altered mental status
- Weakness / Paralysis
- Blindness or other sensory loss
- Aphasia / Dysarthria
- Syncope
- Vertigo / Dizziness
- Vomiting
- Headache
- Seizures
- Respiratory pattern change
- Hypertension / hypotension

Differential
- See Altered Mental Status
- TIA (Transient ischemic attack)
- Seizure
- Todd's Paralysis
- Hypoglycemia
- Stroke
  Thrombotic or Embolic (~85%)
  Hemorrhagic (~15%)
- Tumor
- Trauma
- Dialysis / Renal Failure

Prehospital Stroke Screen

Consistent with Acute Stroke

YES → **Time of Onset Or Time Last Seen Normal < 12 Hours**

YES → **Transport based on:**
STROKE
EMS Triage and Destination Plan
Keep Scene Time to ≤ 10 Minutes
Immediate Notification of Facility

NO → Exit to Appropriate Protocol(s) ***as indicated***

NO →

- Blood Glucose Analysis Procedure
- Altered Mental Status Protocol UP 4 ***if indicated***
- Reperfusion Checklist Procedure
- **B** — 12 Lead ECG Procedure
- **A** — IV Procedure **P** IO Procedure
  Preferably 2 Sites
  Venous Access Blood Draw Procedure ***if applicable***
- **P** — Cardiac Monitor
- Age Appropriate Cardiac Protocol(s) ***if indicated***
- Head Trauma Protocol TB 5
  Multiple Trauma Protocol TB 6 ***if indicated***
- Seizure Protocol UP 13 ***if indicated***

SBP ≥ 220
DBP ≥ 120
after 3 readings
at least 5 minutes apart

YES → Contact Receiving Facility Concerning Treatment of Hypertension

Notify Destination or Contact Medical Control

Universal Protocol Section

Suspected Stroke

Universal Protocol Section

Pearls

- **Recommended Exam: Mental Status, HEENT, Heart, Lungs, Abdomen, Extremities, Neuro**
- **Items in Red Text are key performance measures used in the EMS Acute Stroke Care Toolkit.**
- **Acute Stroke care is evolving rapidly. Time of onset / last seen normal may be changed at any time depending on the capabilities and resources of your hospital based on Stroke: EMS Triage and Destination Plan.**
- **Time of Onset or Last Seen Normal:**
  - **One of the most important items the pre-hospital provider can obtain, of which all treatment decisions are based.**
  - **Be very precise in gathering data to establish the time of onset and report as an actual time (i.e. 13:47 NOT "about 45 minutes ago.")**
  - **Without this information patient may not be able to receive thrombolytics at facility.**
  - **Wake up stroke: Time starts when patient last awake or symptom free.**
- **You are often in the best position to determine the actual Time of Onset while you have family, friends or caretakers available. Often these sources of information may arrive well after you have delivered the patient to the hospital. Delays in decisions due to lack of information may prevent an eligible patient from receiving thrombolytics.**
- **The Reperfusion Checklist should be completed for any suspected stroke patient. With a duration of symptoms of less than** 12 Hours **, scene times should be limited to ≤ 10 minutes, early notification / activation of receiving facility should be performed and transport times should be minimized.**
- **If possible place 2 IV sites.**
- **Blood Draw:**
  - **Many systems utilize EMS venous blood samples. Follow your local policy and procedures.**
- The differential listed on the Altered Mental Status Protocol should also be considered.
- Be alert for airway problems (swallowing difficulty, vomiting/aspiration).
- Hypoglycemia can present as a localized neurologic deficit, especially in the elderly.
- Document the Stroke Screen results in the PCR.
- Agencies may use validated pre-hospital stroke screen of choice.
- **Pediatrics:**
  - Strokes do occur in children, they are slightly more common in ages < 2, in boys, and in African-Americans. Newborn and infant symptoms consist of seizures, extreme sleepiness, and using only one side of the body. Children and teenagers symptoms may consist of severe headaches, vomiting, sleepiness, dizziness, and/or loss of balance or coordination.

Suspected Sepsis

History
- Duration and severity of fever
- Past medical history
- Medications / Recent antibiotics
- Immunocompromised (transplant, HIV, diabetes, cancer)
- Indwelling medical device
- Last acetaminophen or ibuprofen
- Recent Hospital / healthcare facility
- Bedridden or immobile
- Elderly and very young – at risk
- Prosthetic device / indwelling device

Signs and Symptoms
- Warm
- Flushed
- Sweaty
- Chills / Rigors
- Delayed cap refill
- Mental status changes

Associated Symptoms (Helpful to localize source)
- myalgias, cough, chest pain, headache, dysuria, abdominal pain, rash

Differential
- Infections: UTI, Pneumonia, skin/ wound
- Cancer / Tumors / Lymphomas
- Medication or drug reaction
- Connective tissue disease: Arthritis, Vasculitis
- Hyperthyroidism
- Heat Stroke
- Meningitis
- Hypoglycemia/hypothermia
- MI / CVA

Consider: Contact, Droplet, and Airborne Precautions

Temperature Measurement Procedure ***if available***

Fever / Infection Control Protocol UP 10 ***if needed***

Altered Mental Status Protocol UP 4 ***if needed***

B 12 Lead ECG Procedure

A IV Procedure **If indicated** | **P** IO Procedure **If indicated**

P Cardiac Monitor

Sepsis Screen Positive

NO → Exit to Age Appropriate Condition Appropriate Protocol(s)

YES → **SEPSIS ALERT**
Notify Receiving Facility Immediately

A
- Venous Access Blood Draw ***if applicable***
- **Normal Saline 500 mL Bolus**
  Repeat as needed
  Titrate SPB ≥ 90 mmHg
  MAP > 65 mmHg
  Maximum 2 L

  Peds: 20 mL/kg IV / IO
  Repeat to titrate
  Age Appropriate
  SBP ≥ 70 + 2 x Age
  Maximum 60 mL/kg

Age Appropriate Hypotension / Shock Protocol AM 5 / PM 3

Adult SIRS Criteria
Temperature
≥ 100.4° F (38° C)
Or
≤ 96.8° F (36° C)
AND
Any 1 of the following:
HR > 90
RR > 20
EtCO < 25 mmHg

Adult qSOFA Criteria
SBP ≤ 100 mmHG
RR ≥ 22
AMS or new mental status change

Pediatrics SIRS Criteria
Temperature
Same as adult

AND
Heart Rate
1 month – 1 year > 180
2 – 5 years > 140
6 – 12 years > 130
13 – 18 years > 120

MAP (Mean Arterial Pressure)

$$\frac{SBP + 2(DBP)}{3}$$

Monitor usually calculates this value on screen

Notify Destination or Contact Medical Control

Universal Protocol Section

Suspected Sepsis

Universal Protocol Section

- **Pearls**
- **Recommended Exam: Mental Status, Skin, HEENT, Neck, Heart, Lungs, Abdomen, Back, Extremities, Neuro**
- **Recommended Exam Pediatrics: In childhood, physical assessment reveals important clues for sepsis. Look for mental status abnormalities such as anxiety, restlessness, agitation, irritability, confusion, or lethargy. Cardiovascular findings to look for include cool extremities, capillary refill >3 seconds, or mottled skin.**
- **Sepsis is a life threatening condition where the body's immune response to infection injures its own tissues and organs.**
- **Severe sepsis is a suspected infection and 2 or more SIRS criteria (or qSOFA) with organ dysfunction such as AMS or hypotension.**
- **Septic shock is severe sepsis and poor perfusion unimproved after fluid bolus.**
- **Agencies administering antibiotics should inquire about drug allergies specific to antibiotics or family of antibiotics.**
- **Following each fluid bolus, assess for pulmonary edema. Consider administration of agency specific vasopressor.**
- **Supplemental oxygen should be given and titrated to oxygenation saturation ≥ 94%.**
- **EKG should be obtained with suspected sepsis, but should not delay care in order to obtain.**
- **Abnormally low temperatures increase mortality and found often in geriatric patients.**
- Quantitative waveform capnography can be a reliable surrogate for lactate monitoring in detecting metabolic distress in sepsis patients. $EtCO_2$ < 25 mm Hg are associated with serum lactate levels > 4 mmol/L.
- Patients with a history of liver failure should not receive acetaminophen.
- **Droplet precautions:**
  - Include standard PPE plus a standard surgical mask for providers who accompany patients in the back of the ambulance and a surgical mask or NRB O2 mask for the patient.
  - This level of precaution should be utilized when influenza, meningitis, mumps, streptococcal pharyngitis, and other illnesses spread via large particle droplets are suspected.
  - A patient with a potentially infectious rash should be treated with droplet precautions.
- **Airborne precautions:**
  - Include standard PPE plus utilization of a gown, change of gloves after every patient contact, and strict hand washing precautions.
  - This level of precaution is utilized when multi-drug resistant organisms (e.g. MRSA), scabies, or zoster (shingles), or other illnesses spread by contact are suspected.
- **All-hazards precautions:**
  - Include standard PPE plus airborne precautions plus contact precautions.
  - This level of precaution is utilized during the initial phases of an outbreak when the etiology of the infection is unknown or when the causative agent is found to be highly contagious (e.g. SARS).
- All patients should have drug allergies documented prior to administering pain medications.
- Allergies to NSAIDs (non-steroidal anti-inflammatory medications) are a contraindication to Ibuprofen.
- Agency Medical Director may require contact of medical control prior to EMT / MR administering any medication.
- **Sepsis Screen:**
  - Agencies may use Adult / Pediatric Systemic Inflammatory Response Syndrome (SIRS) criteria or quickSOFA (qSOFA) criteria.
  - Receiving facility should be involved in determining Sepsis Screen utilized by EMS.

Syncope

History
- Cardiac history, stroke, seizure
- Occult blood loss (GI, ectopic)
- Females: LMP, vaginal bleeding
- Fluid loss: nausea, vomiting, diarrhea
- Past medical history
- Medications

Signs and Symptoms
- Loss of consciousness with recovery
- Lightheadedness, dizziness
- Palpitations, slow or rapid pulse
- Pulse irregularity
- Decreased blood pressure

Differential
- Vasovagal
- Orthostatic hypotension
- Cardiac syncope
- Micturition / Defecation syncope
- Psychiatric
- Stroke
- Hypoglycemia
- Seizure
- Shock (see Shock Protocol)
- Toxicological (Alcohol)
- Medication effect (hypertension)
- PE
- AAA

Age Appropriate Airway Protocol(s) AR 1, 2, 3, 5, 6 ***if indicated***

Blood Glucose Analysis Procedure

B 12 Lead ECG Procedure

I IV Procedure **P** IO Procedure

P Cardiac Monitor

Altered Mental Status Protocol UP 4 ***if indicated***

Age Appropriate Cardiac Protocol(s) ***if indicated***

Age Appropriate Hypotension / Shock Protocol AM 5 / PM 3 ***if indicated***

Multiple Trauma Protocol TB 6 Spinal Motion Restriction Procedure / Protocol TB 8 ***if indicated***

Serious Signs / Symptoms Hypotension, poor perfusion, shock

YES →

A

IV Procedure **P** IO Procedure

Consider 2 Large Bore Sites

Normal Saline 500 mL Bolus
Repeat as needed
Titrate SPB ≥ 90 mmHg
Maximum 2 L
Peds: 20 mL/kg IV / IO
Repeat as needed
Titrate to Age Appropriate
SBP ≥ 70 + 2 x Age
Maximum 60 mL/kg

Exit to Age Appropriate Condition Appropriate Protocol(s)

NO

Notify Destination or Contact Medical Control

Universal Protocol Section

Syncope

Universal Protocol Section

Pearls

- **Recommended Exam: Mental Status, Skin, HEENT, Heart, Lungs, Abdomen, Back, Extremities, Neuro**
- **Syncope is both loss of consciousness and loss of postural tone. Symptoms preceding the event are important in determining etiology.**
- **Syncope often is due to a benign process but can be an indication of serious underlying disease in both the adult and pediatric patient.**
- **Often patients with syncope are found normal on EMS evaluation. In general patients experiencing syncope require cardiac monitoring and emergency department evaluation.**
- **Differential should remain wide and include:**

Cardiac arrhythmia	Neurological problem	Choking	Pulmonary embolism
Hemorrhage	Stroke	Respiratory	Hypo or Hyperglycemia
GI Hemorrhage	Seizure	Sepsis	

- **High-risk patients:**

Age ≥ 60	Syncope with exertion
History of CHF	Syncope with chest pain
Abnormal ECG	Syncope with dyspnea

- **Age specific blood pressure 0 – 28 days > 60 mmHg, 1 month - 1 year > 70 mmHg, 1 - 10 years > 70 + (2 x age) mmHg and 11 years and older > 90 mmHg.**
- **Abdominal / back pain in women of childbearing age should be treated as pregnancy related until proven otherwise.**
- **The diagnosis of abdominal aneurysm should be considered with abdominal pain, with or without back and / or lower extremity pain or diminished pulses, especially in patients over 50 and / or patients with shock/ poor perfusion. Notify receiving facility early with suspected abdominal aneurysm.**
- **Consider cardiac etiology in patients > 50, diabetics and / or women especially with upper abdominal complaints.**
- **Heart Rate: One of the first clinical signs of dehydration, almost always increased heart rate, tachycardia increases as dehydration becomes more severe, very unlikely to be significantly dehydrated if heart rate is close to normal.**
- Syncope with no preceding symptoms or event may be associated with arrhythmia.
- Assess for signs and symptoms of trauma if associated or questionable fall with syncope.
- Consider dysrhythmias, GI bleed, ectopic pregnancy, and seizure as possible causes of syncope.
- These patients should be transported. Patients who experience syncope associated with headache, neck pain, chest pain, abdominal pain, back pain, dyspnea, or dyspnea on exertion need prompt medical evaluation.
- More than 25% of geriatric syncope is cardiac dysrhythmia based.