

Bites and Envenomations

History


- Type of bite / sting
- Description or bring creature / photo with patient for identification
- Time, location, size of bite / sting
- Previous reaction to bite / sting
- Domestic vs. Wild
- Tetanus and Rabies risk
- Immunocompromised patient

Signs and Symptoms

- Rash, skin break, wound
- Pain, soft tissue swelling, redness
- Blood oozing from the bite wound
- Evidence of infection
- Shortness of breath, wheezing
- Allergic reaction, hives, itching
- Hypotension or shock

Differential

- Animal bite
- Human bite
- Snake bite (poisonous)
- Spider bite (poisonous)
- Insect sting / bite (bee, wasp, ant, tick)
- Infection risk
- Rabies risk
- Tetanus risk



Bites and Envenomations

Pearls

- **Recommended Exam: Mental Status, Skin, Extremities (Location of injury), and a complete Neck, Lung, Heart, Abdomen, Back, and Neuro exam if systemic effects are noted**
- Human bites have higher infection rates than animal bites due to normal mouth bacteria.
- Carnivore bites are much more likely to become infected and all have risk of Rabies exposure.
- Cat bites may progress to infection rapidly due to a specific bacteria (*Pasteurella multocida*).
- Poisonous snakes in this area are generally of the pit viper family: rattlesnake and copperhead.
- Coral snake bites are rare: Very little pain but very toxic. "Red on yellow - kill a fellow, red on black - venom lack."
- Amount of envenomation is variable, generally worse with larger snakes and early in spring.
- If no pain or swelling, envenomation is unlikely. About 25 % of snake bites are "dry" bites.
- Black Widow spider bites tend to be minimally painful, but over a few hours, muscular pain and severe abdominal pain may develop (spider is black with red hourglass on belly).
- Brown Recluse spider bites are minimally painful to painless. Little reaction is noted initially but tissue necrosis at the site of the bite develops over the next few days (brown spider with fiddle shape on back).
- Evidence of infection: swelling, redness, drainage, fever, red streaks proximal to wound.
- Immunocompromised patients are at an increased risk for infection: diabetes, chemotherapy, transplant patients.
- Consider contacting the North Carolina Poison Control Center for guidance (1-800-84-TOXIN).

Carbon Monoxide / Cyanide

History


- Smoke inhalation
- Ingestion of cyanide
- Eating large quantity of fruit pits
- Industrial exposure
- Trauma
- Reason: Suicide, criminal, accidental
- Past Medical History
- Time / Duration of exposure

Signs and Symptoms

- AMS
- Malaise, weakness, flu like illness
- Dyspnea
- GI Symptoms; N/V; cramping
- Dizziness
- Seizures
- Syncope
- Reddened skin
- Chest pain

Differential

- Diabetic related
- Infection
- MI
- Anaphylaxis
- Renal failure / dialysis problem
- Head injury / trauma
- Co-ingestant or exposures



Pearls

- **Recommended exam: Neuro, Skin, Heart, Lungs, Abdomen, Extremities**
- **Scene safety is priority.**
- Consider CO and Cyanide with any product of combustion
- Normal environmental CO level does not exclude CO poisoning.
- Symptoms present with lower CO levels in pregnancy, children and the elderly.
- Continue high flow oxygen regardless of pulse ox readings.

Drowning / Submersion Injury

History


- Submersion in water regardless of depth
- Possible history of trauma ie: diving board
- Duration of immersion
- Temperature of water or possibility of hypothermia
- Degree of water contamination

Signs and Symptoms

- Unresponsive
- Mental status changes
- Decreased or absent vital signs
- Vomiting
- Coughing, Wheezing, Rales, Rhonci, Stridor
- Apnea

Differential

- Trauma
- Pre-existing medical problem
- Pressure injury (diving)
- Barotrauma
- Decompression sickness
- Post-immersion syndrome



Pearls

- **Recommended Exam: Trauma Survey, Head, Neck, Chest, Abdomen, Pelvis, Back, Extremities, Skin, Neuro**
- **Ensure scene safety. Drowning is a leading cause of death among would-be rescuers.**
- **Allow appropriately trained and certified rescuers to remove victims from areas of danger.**
- **With cold water no time limit -- resuscitate all. These patients have an increased chance of survival.**
- Have a high index of suspicion for possible spinal injuries
- Hypothermia is often associated with drowning and submersion injuries.
- All victims should be transported for evaluation due to potential for worsening over the next several hours.
- With pressure injuries (decompression / barotrauma), consider transport to or availability of a hyperbaric chamber.

Hyperthermia

History


- Age, very young and old
- Exposure to increased temperatures and / or humidity
- Past medical history / Medications
- Time and duration of exposure
- Poor PO intake, extreme exertion
- Fatigue and / or muscle cramping

Signs and Symptoms

- Altered mental status / coma
- Hot, dry or sweaty skin
- Hypotension or shock
- Seizures
- Nausea

Differential

- Fever (Infection)
- Dehydration
- Medications
- Hyperthyroidism (Storm)
- Delirium tremens (DT's)
- Heat cramps, exhaustion, stroke
- CNS lesions or tumors



Hyperthermia

Pearls

- **Recommended Exam: Mental Status, Skin, HEENT, Heart, Lungs, Neuro**
- Extremes of age are more prone to heat emergencies (i.e. young and old). Obtain and document patient temperature if able.
- Predisposed by use of: tricyclic antidepressants, phenothiazines, anticholinergic medications, and alcohol.
- Cocaine, Amphetamines, and Salicylates may elevate body temperatures.
- Sweating generally disappears as body temperature rises above 104° F (40° C).
- Intense shivering may occur as patient is cooled.
- **Heat Cramps** consists of benign muscle cramping 2° to dehydration and is not associated with an elevated temperature.
- **Heat Exhaustion** consists of dehydration, salt depletion, dizziness, fever, mental status changes, headache, cramping, nausea and vomiting. Vital signs usually consist of tachycardia, hypotension, and an elevated temperature.
- **Heat Stroke** consists of dehydration, tachycardia, hypotension, temperature >104° F (40° C), and an altered mental status.

Hypothermia / Frostbite

History


- Age, very young and old
- Exposure to decreased temperatures but may occur in normal temperatures
- Past medical history / Medications
- Drug use: Alcohol, barbituates
- Infections / Sepsis
- Length of exposure / Wetness / Wind chill

Signs and Symptoms

- Altered mental status / coma
- Cold, clammy
- Shivering
- Extremity pain or sensory abnormality
- Bradycardia
- Hypotension or shock

Differential

- Sepsis
- Environmental exposure
- Hypoglycemia
- CNS dysfunction
 - Stroke
 - Head injury
 - Spinal cord injury



Hypothermia / Frostbite

Pearls

- **Recommended Exam: Mental Status, Heart, Lungs, Abdomen, Extremities, Neuro**
- **NO PATIENT IS DEAD UNTIL WARM AND DEAD (Body temperature \geq 93.2 degrees F, 32 degrees C.)**
- **Hypothermia categories:**
 - Mild 90 – 95 degrees F (32 – 35 degrees C)
 - Moderate 82 – 90 degrees F (28 – 32 degrees C)
 - Severe < 82 degrees F (< 28 degrees C)
- **Mechanisms of hypothermia:**
 - Radiation: Heat loss to surrounding objects via infrared energy (60 % of most heat loss.)
 - Convection: Direct transfer of heat to the surrounding air.
 - Conduction: Direct transfer of heat to direct contact with cooler objects (important in submersion.)
 - Evaporation: Vaporization of water from sweat or other body water losses.
- Contributing factors of hypothermia: Extremes of age, malnutrition, alcohol or other drug use.
- If the temperature is unable to be measured, treat the patient based on the suspected temperature.
- **CPR:**
 - Severe hypothermia may cause cardiac instability and rough handling of the patient theoretically can cause ventricular fibrillation. This has not been demonstrated or confirmed by current evidence.
 - Intubation and CPR techniques should not be with-held due to this concern.
 - Intubation can cause ventricular fibrillation so it should be done gently by most experienced person.
 - Below 86 degrees F (30 degrees C) antiarrhythmics may not work and if given should be given at reduced intervals. Contact medical control for direction. Epinephrine / Vasopressin can be administered. Below 86 degrees F (30 degrees) pacing should not be done
 - Consider withholding CPR if patient has organized rhythm or has other signs of life. Contact Medical Control.
 - If the patient is below 86 degrees F (30 degree C) then defibrillate 1 time if defibrillation is required. Deferring further attempts until more warming occurs is controversial. Contact medical control for direction.
 - Hypothermia may produce severe bradycardia so take at least 45 second to palpate a pulse.
- Hot packs can be activated and placed in the armpit and groin area if available. Care should be taken not to place the packs directly against the patient's skin.

Marine Envenomations / Injury

History


- Type of bite / sting
- Identification of organism
- Previous reaction to marine organism
- Immunocompromised
- Household pet

Signs and Symptoms

- Intense localized pain
- Increased oral secretions
- Nausea / vomiting
- Abdominal cramping
- Allergic reaction / anaphylaxis

Differential

- Jellyfish sting
- Sea Urchin sting
- Sting ray barb
- Coral sting
- Swimmers itch
- Cone Shell sting
- Fish bite
- Lion Fish sting



Marine Envenomations / Injury

Pearls

- Ensure your safety: Avoid the organism or fragments of the organism as they may impart further sting / injury.
- Patients can suffer cardiovascular collapse from both the venom and / or anaphylaxis even in seemingly minor envenomations.
- Sea creature stings and bites impart moderate to severe pain.
- Arrest the envenomation by inactivation of the venom as appropriate.
- Ensure good wound care, immobilization and pain control.

WMD-Nerve Agent Protocol

History


- Exposure to chemical, biologic, radiologic, or nuclear hazard
- Potential exposure to unknown substance/hazard

Signs and Symptoms

- Salivation
- Lacrimation
- Urination; increased, loss of control
- Defecation / Diarrhea
- GI Upset; Abdominal pain / cramping
- Emesis
- Muscle Twitching
- Seizure Activity
- Respiratory Arrest

Differential

- Nerve agent exposure (e.g., VX, Sarin, Soman, etc.)
- Organophosphate exposure (pesticide)
- Vesicant exposure (e.g., Mustard Gas, etc.)
- Respiratory Irritant Exposure (e.g., Hydrogen Sulfide, Ammonia, Chlorine, etc.)



Pearls

- **Recommended Exam:** Mental Status, Skin, HEENT, Heart, Lungs, Gastrointestinal, Neuro
- **Follow local HAZMAT protocols for decontamination and use of personal protective equipment.**
- In the face of a bona fide attack, begin with 1 Nerve Agent Kit for patients less than 7 years of age, 2 Nerve Agent Kits from 8 to 14 years of age, and 3 Nerve Agent Kits for patients 15 years of age and over.
- If Triage/MCI issues exhaust supply of Nerve Agent Kits, use pediatric atropines (if available). Use the 0.5 mg dose if patient is less than 40 pounds (18 kg), 1 mg dose if patient weighs between 40 to 90 pounds (18 to 40 kg), and 2 mg dose for patients greater than 90 pounds (>40 kg).
- Each Nerve Agent Kit contains 600 mg of Pralidoxime (2-PAM) and 2 mg of Atropine.
- **Seizure Activity:** Any benzodiazepine by any route is acceptable.
- For patients with major symptoms, there is no limit for atropine dosing.
- Carefully evaluate patients to ensure they not from exposure to another agent (e.g., narcotics, vesicants, etc.)
- The main symptom that the atropine addresses is excessive secretions so atropine should be given until salivation improves.
- EMS personnel, public safety officers and Medical Responders / EMT-B may carry, self-administer or administer to a patient atropine / pralidoxime by protocol. Agency medical director may require Contact of Medical Control prior to administration.